arXiv:0907.0184v1 [hep-ph] 1 Jul 2009

Open charm hadroproduction and the intrinsic charm content of the proton

Bernd A. Kniehl

II. Institut für Theoretische Physik, Universität Hamburg,
Luruper Chaussee 149, 22761 Hamburg, Germany

We advocate charmed-hadron inclusive hadroproduction as a laboratory to probe intrinsic charm (IC) inside the colliding hadrons. Working at next-to-leading order in the general-mass variable-flavor-number scheme endowed with non-perturbative fragmentation functions recently extracted from a global fit to e^+e^- annihilation data from KEKB, CESR, and LEP1, we first assess the sensitivity of Tevatron data of D^0, D^+, and D^{*+} inclusive production to the IC parameterizations provided by Pumplin *et al.* We then argue that similar data from pp collisions at RHIC would have the potential to discriminate between different IC models provided they reach out to sufficiently large values of transverse momentum.

1 Introduction

The general-mass variable-flavor-number scheme (GM-VFNS) provides a rigorous theoretical framework for the theoretical description of the inclusive production of single heavy-flavored hadrons, combining the fixed-flavor-number scheme (FFNS) and zero-mass variable-flavor-number scheme (ZM-FVNS), which are valid in complementary kinematic regions, in a unified approach that enjoys the virtues of both schemes and, at the same time, is bare of their flaws. Specifically, it resums large logarithms by the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) evolution of non-perturbative fragmentation functions (FFs), guarantees the universality of the latter as in the ZM-VFNS, and simultaneously retains the mass-dependent terms of the FFNS without additional theoretical assumptions. It was elaborated at next-to-leading order (NLO) for e^+e^- annihilation [2] photo- [3] and hadroproduction [4, 5]. In this presentation [1], we report recent progress in the theoretical description of D-meson hadroproduction and argue that this process provides a useful probe of IC in the proton [6]. In Sec. 2, we present updated predictions for the transverse-momentum (p_T) distributions of D^0, D^+, and D^{*+} mesons [6] using our new FF sets [2]. In Sec. 3, we briefly review the constraints on IC obtained through an extension [7] of the CTEQ6.5 global analysis [8] endowed with three representative IC models. In Sec. 4, we investigate, for D^0 hadroproduction at the Tevatron and at RHIC, the shifts in cross section produced by these IC implementations. Our conclusions are summarized in Sec. 5.

2 D-meson hadroproduction revisited

Our previous NLO predictions for inclusive D-meson hadroproduction [5] were evaluated with the FF sets of Ref. [9]. These and previous FF sets [10] had been determined through fits to LEP1 data [11] in the ZM-VFNS. These determinations have recently been updated and improved [2] by also including KEKB and CESR [12] data, switching to the GM-VFNS, and also incorporating hadron-mass corrections. They greatly improve the agreement with the p_T distributions of D^0, D^+, and D^{*+} mesons measured at the Tevatron [13], as may be seen in Fig. 1(a)–(c) and (d)–(f) on an absolute scale and in the data-over-theory representation, respectively.



(a) (b) (c)

(d) (e) (f)

Figure 1: The solid lines represent the default predictions, and the dashed lines indicate the theoretical uncertainty.

3 Intrinsic charm from global analysis

In Ref. [7], the CTEQ6.5 global analysis [8] was repeated allowing for IC according to the Brodsky-Hoyer-Peterson-Sakai (BHPS) [14], meson-cloud [15], and sea-like [7] models, with a modest and a marginal value of $\langle x \rangle_{c+\overline{c}}$ in each case. The x distributions at $\mu = 2$ GeV of the resulting c and $\overline{c}$ parton distribution functions (PDFs) are shown in Fig. 2(a)–(c) for the BHPS, meson-cloud, and sea-like models, respectively.

4 Probing intrinsic charm at Tevatron and RHIC

In the GM-VFNS, D-meson hadroproduction receives sizeable contributions from partonic subprocesses with incoming c or $\overline{c}$ quarks [4, 5] and is thus sensitive to their PDFs and IC-induced enhancements in them. In Fig. 3(a), the relative enhancements due to the six



(a) (b) (c)

Figure 2: The lower and upper dashed lines correspond to IC with modest and marginal $\langle x\rangle_{c+\bar{c}}$ values, respectively, and the solid ones to zero IC.

IC implementations of Ref. [7] are compared with the D^0 data of Ref. [13]. We observe that the two large-p_T data points tend to disfavour the sea-like IC with marginal value of $\langle x\rangle_{c+\bar{c}}$, although a firm statement would be premature. On the other hand, the effects due the IC of the BHPS or meson-cloud models are too feeble to be resolved by the presently published CDF data.

Since IC typically receives a large fraction x of momentum from the parent hadron, especially in the BHPS and meson-cloud models, and the kinematical upper bound on x scales with $x_T = 2p_T/\sqrt{s}$, the sensitivity to IC may be enhanced by increasing p_T or lowering the cm. energy $\sqrt{s}$. Keeping in mind that the CDF analysis of Ref. [13] is merely based on 5.8 pb^{-1} of data recorded in February and March 2002 and that, since then, the integrated luminosity of run II has increased by more than a factor of 1000, exceeding 6 fb^{-1} as it does, significantly more precise data at larger p_T values are expected to come, with an increased sensitivity to IC, as may be seen from Fig. 3(b). A hadron collider with smaller c.m. energy is in operation, namely RHIC with presently $\sqrt{s} = 200$ GeV, where the p_T distribution of $pp \to X_c + X$ could be measured. So far, only the STAR Collaboration has published D-meson production data, namely for $d\mathrm{Au} \to D^0 + X$ with $p_T < 2.5$ GeV [16]. In Fig. 3(c), we observe significant enhancements due to IC in pp collisions at $\sqrt{s} = 200$ GeV, by up to 75% at $p_T = 10$ GeV. The high-energy pp mode of RHIC, with $\sqrt{s} = 500$ GeV, will accrue more luminosity, perhaps 200–400 pb^{-1}. However, this will happen at the expense of lowering the accessible x values and, thus, of reducing the relative IC effects on the cross sections, especially for the BHPS and meson-cloud models, as is evident from Fig. 3(d).

5 Conclusions

In the case of the Tevatron, we found that the BHPS and meson-cloud models yield insignificant deviations from the zero-IC predictions, while the shift produced by the sea-like model is comparable to the experimental error and tends to worsen the agreement between theory and experiment. However, the experimental errors in Ref. [13] are still too sizeable to rule out this model as implemented in Ref. [7]. This is likely to change once the full data sample of run II is exploited.

As for pp collisions with $\sqrt{s} = 200$ GeV at RHIC, all three IC models predict sizeable enhancements, by up to 75% at $p_T = 10$ GeV. In a future high-energy pp-collision mode of

RHIC, with $\sqrt{s} = 500$ GeV, which is to accrue more luminosity, the cross sections would be increased, while the relative shifts due to IC would be considerably smaller for the BHPS and meson-cloud models.



(a)

(b)

(c)

(d)

Figure 3: The solid (dashed), densely dotted (dot-dashed), and scarcely dotted (dotted) lines correspond to the BHPS [14], meson-cloud [15], and sea-like [7] models with modest (marginal) values of $\langle x \rangle_{c+\bar{c}}$.

Acknowledgments

The author thanks G. Kramer, I. Schienbein, and H. Spiesberger for the collaboration on the work presented here. This work was supported in part by the Bundesministerium für Bildung und Forschung BMBF through Grant No. 05 HT6GUA, by the Deutsche Forschungsgemeinschaft DFG through Grant No. KN 365/7–1, and by the Helmholtz-Gemeinschaft Deutscher Forschungszentren HGF through Grant No. HA–101.

References

[1] Slides:
`http://indico.cern.ch/contributionDisplay.py?contribId=138&sessionId=5&confId=53294`

[2] T. Kneesch, B. A. Kniehl, G. Kramer, and I. Schienbein, Nucl. Phys. **B799**, 34 (2008) [arXiv:0712.0481 [hep-ph]]; B. A. Kniehl, in *Proceedings of the XVI International Workshop on Deep-Inelastic Scattering and Related Subjects (DIS 2008)*, London, England, 2008, edited by R. Devenish and J. Ferrando (Science Wise Publishing, Amsterdam, 2008), `http://dx.doi.org/10.3360/dis.2008.195` [arXiv:0807.2215 [hep-ph]].

[3] G. Kramer and H. Spiesberger, Eur. Phys. J. C **22**, 289 (2001) [arXiv:hep-ph/0109167]; Eur. Phys. J. C **28**, 495 (2003) [arXiv:hep-ph/0302081]; Eur. Phys. J. C **38**, 309 (2004) [arXiv:hep-ph/0311062]; B. A. Kniehl, G. Kramer, I. Schienbein, and H. Spiesberger, Eur. Phys. J. C **62**, 365 (2009) [arXiv:0902.3166 [hep-ph]].

[4] B. A. Kniehl, G. Kramer, I. Schienbein, and H. Spiesberger, Phys. Rev. D **71**, 014018 (2005) [arXiv:hep-ph/0410289]; Eur. Phys. J. C **41**, 199 (2005) [arXiv:hep-ph/0502194].

[5] B. A. Kniehl, G. Kramer, I. Schienbein, and H. Spiesberger, AIP Conf. Proc. **792**, 867 (2005) [arXiv:hep-ph/0507068]; Phys. Rev. Lett. **96**, 012001 (2006) [arXiv:hep-ph/0508129]; B. A. Kniehl, in *Proceedings of the 14th International Workshop on Deep Inelastic Scattering (DIS 2006)*, Tsukuba, Japan, 2006, edited by M. Kuze, K. Nagano, and K. Tokushuku (World Scientific, Singapore, 2007) p. 573 [arXiv:hep-ph/0608122].

[6] B. A. Kniehl, G. Kramer, I. Schienbein, and H. Spiesberger, Phys. Rev. D **79**, 094009 (2009) [arXiv:0901.4130 [hep-ph]].

[7] J. Pumplin, H. L. Lai, and W. K. Tung, Phys. Rev. D **75**, 054029 (2007) [arXiv:hep-ph/0701220].

[8] CTEQ Collaboration, W. K. Tung, H. L. Lai, A. Belyaev, J. Pumplin, D. Stump, and C. P. Yuan, JHEP **0702**, 053 (2007) [arXiv:hep-ph/0611254].

[9] B. A. Kniehl and G. Kramer, Phys. Rev. D **74**, 037502 (2006) [arXiv:hep-ph/0607306].

[10] J. Binnewies, B. A. Kniehl, and G. Kramer, Phys. Rev. D **58**, 014014 (1998) [arXiv:hep-ph/9712482]; B. A. Kniehl and G. Kramer, Phys. Rev. D **71**, 094013 (2005) [arXiv:hep-ph/0504058].

[11] OPAL Collaboration, G. Alexander *et al.*, Z. Phys. C **72**, 1 (1996); OPAL Collaboration, K. Ackerstaff *et al.*, Eur. Phys. J. C **1**, 439 (1998) [arXiv:hep-ex/9708021]; ALEPH Collaboration, R. Barate *et al.*, Eur. Phys. J. C **16**, 597 (2000) [arXiv:hep-ex/9909032].

[12] CLEO Collaboration, M. Artuso *et al.*, Phys. Rev. D **70**, 112001 (2004) [arXiv:hep-ex/0402040]; Belle Collaboration, R. Seuster *et al.*, Phys. Rev. D **73**, 032002 (2006) [arXiv:hep-ex/0506068].

[13] CDF Collaboration, D. E. Acosta *et al.*, Phys. Rev. Lett. **91**, 241804 (2003) [arXiv:hep-ex/0307080].

[14] S. J. Brodsky, P. Hoyer, C. Peterson, and N. Sakai, Phys. Lett. B **93**, 451 (1980); S. J. Brodsky, C. Peterson, and N. Sakai, Phys. Rev. D **23**, 2745 (1981).

[15] F. S. Navarra, M. Nielsen, C. A. A. Nunes, and M. Teixeira, Phys. Rev. D **54**, 842 (1996) [arXiv:hep-ph/9504388]; S. Paiva, M. Nielsen, F. S. Navarra, F. O. Duraes, and L. L. Barz, Mod. Phys. Lett. A **13**, 2715 (1998) [arXiv:hep-ph/9610310]; W. Melnitchouk and A. W. Thomas, Phys. Lett. B **414**, 134 (1997) [arXiv:hep-ph/9707387]; F. M. Steffens, W. Melnitchouk, and A. W. Thomas, Eur. Phys. J. C **11**, 673 (1999) [arXiv:hep-ph/9903441].

[16] STAR Collaboration, J. Adams *et al.*, Phys. Rev. Lett. **94**, 062301 (2005) [arXiv:nucl-ex/0407006].